

## Modification of Carmone, Kara & Maxwell Heuristic Identification of Noisy Variables (HINoV)

### Algorithm for metric data (see Carmone, Kara and Maxwell [1999])

- Step 1.** Data matrix containing  $m$  normalized variables measured on metric scale (ratio, interval) and  $n$  objects ( $i = 1, \dots, n$ ;  $j = 1, \dots, m$ ) is a starting point.
- Step 2.** Cluster, via `kmeans` method, the observed data separately for each  $j$ -th variable for a given number of cluster  $u$ . It is possible to use clustering methods based on distance matrix (`pam` or any hierarchical agglomerative method: `single`, `complete`, `average`, `mcquitty`, `median`, `centroid`, `ward.D`, `ward.D2`).
- Step 3.** Calculate adjusted Rand indices  $R_{jl}$  ( $j, l = 1, \dots, m$ ) for partitions formed from all distinct pairs of the  $m$  variables ( $j \neq l$ ). Due to fact that adjusted Rand (Rand) index is symmetrical we need to calculate  $m(m - 1)/2$  values.
- Step 4.** Construct  $m \times m$  adjusted Rand matrix (`parim`). Sum rows (or columns) for each  $j$ -th variable  $R_{j\bullet} = \sum_{l=1}^m R_{jl}$  (`topri`):

$$\begin{array}{c} \begin{bmatrix} M_1 \\ M_2 \\ \vdots \\ M_j \\ \vdots \\ M_m \end{bmatrix} \end{array} \begin{array}{c} \text{parim} \\ \begin{bmatrix} R_{12} & \dots & R_{1l} & \dots & R_{1m} \\ R_{21} & & \dots & R_{2l} & \dots & R_{2m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ R_{j1} & R_{j2} & \dots & R_{jl} & \dots & R_{jm} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ R_{m1} & R_{m2} & \dots & R_{ml} & \dots & \end{bmatrix} \end{array} \begin{array}{c} \text{topri} \\ \begin{bmatrix} R_{1\bullet} \\ R_{2\bullet} \\ \vdots \\ R_{j\bullet} \\ \vdots \\ R_{m\bullet} \end{bmatrix} \end{array}$$

- Step 5.** Rank `topri` values  $R_{1\bullet}, R_{2\bullet}, \dots, R_{m\bullet}$  in decreasing order (`stopri`) and plot the scree diagram. The size of the `topri` values indicate the contribution of that variable to the cluster structure. A scree diagram identifies sharp changes in `topri` values. Relatively low-valued `topri` variables (the noisy variables) are identified and eliminated from further analysis (say  $h$  variables).
- Step 6.** Run cluster analysis (based on the same classification method) with the selected  $m - h$  variables.

Modification of Carmone, Kara & Maxwell Heuristic Identification of Noisy Variables (HINoV) method for nonmetric data<sup>1</sup> differs in steps 1, 2, and 6 (see Walesiak [2005], Walesiak and Dudek [2008]):

- Step 1.** Data matrix  $[x_{ij}]$  containing  $m$  ordinal and/or nominal variables and  $n$  objects is a starting point.
- Step 2.** For each  $j$ -th variable we receive natural clusters, where number of clusters equals number of categories for that variable (for instance five for Likert scale or seven for semantic differential scale).
- Step 6.** Run cluster analysis with one of clustering methods based on distance appropriate to nonmetric data (GDM2 for ordinal data – see Jajuga, Walesiak & Bak [2003]; Sokal and Michener distance for nominal data) with the selected  $m - h$  variables.

---

<sup>1</sup> For nonmetric variables (ordinal, nominal) contain not to many categories (for nonmetric variables where number of objects is much more than number of categories).

## References

- Carmone F.J., Kara A., Maxwell S. (1999), *HINoV: a new method to improve market segment definition by identifying noisy variables*, "Journal of Marketing Research", November, vol. 36, 501-509.
- Hubert L.J., Arabie P. (1985), *Comparing partitions*, "Journal of Classification", no. 1, 193-218.
- Jajuga, K., Walesiak, M., Bąk, A. (2003): *On the General Distance Measure*, In: M. Schwaiger, and O. Opitz (Eds.), *Exploratory Data Analysis in Empirical Research*, Springer-Verlag, Berlin, Heidelberg, 104-109. DOI: [https://doi.org/10.1007/978-3-642-55721-7\\_12](https://doi.org/10.1007/978-3-642-55721-7_12)
- Rand W.M. (1971), *Objective criteria for the evaluation of clustering methods*, "Journal of the American Statistical Association", no. 336, 846-850.
- Walesiak M. (2005), *Variable selection for cluster analysis – approaches, problems, methods*, Plenary Session of the Committee on Statistics and Econometrics of the Polish Academy of Sciences, 15 March, Wrocław.
- Walesiak, M., Dudek, A. (2008), *Identification of noisy variables for nonmetric and symbolic data in cluster analysis*, In: C. Preisach, H. Burkhardt, L. Schmidt-Thieme, R. Decker (red.), *Data analysis, machine learning and applications*, Springer-Verlag, Berlin, Heidelberg, 85-92. DOI: [https://doi.org/10.1007/978-3-540-78246-9\\_11](https://doi.org/10.1007/978-3-540-78246-9_11)